

## **SUPPLEMENTAL MATERIAL**

### **Maternal Urinary Bisphenol A during Pregnancy and Maternal and Neonatal Thyroid Function in the CHAMACOS Study**

Jonathan Chevrier<sup>1</sup>, Robert B. Gunier<sup>1</sup>, Asa Bradman<sup>1</sup>, Nina T. Holland<sup>1</sup>, Antonia M. Calafat<sup>2</sup>, Brenda Eskenazi<sup>1</sup> and Kim G. Harley<sup>1</sup>

<sup>1</sup>Center for Children's Environmental Health Research, School of Public Health, University of California, Berkeley, Berkeley, California, USA.

<sup>2</sup>Division for Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA.

**Supplemental Material, Table S1.** Urinary bisphenol A concentrations ( $\mu\text{g/g}$  creatinine) during pregnancy in CHAMACOS study participants in samples included in analyses of maternal (n=335) and neonatal (n=364) serum thyroid hormone levels.

Timing of Measurement	N	Percentiles			Range	Geometric Mean (GSD <sup>a</sup> )	Detection Frequency
		25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>			
<b>Maternal TH<sup>b</sup> analyses</b>							
Closest measurement	335	0.7	1.1	1.9	<LOD <sup>c</sup> – 37	1.2 (2.3)	82%
Farthest measurement	335	0.7	1.1	1.8	<LOD – 27	1.1 (2.3)	81%
Pregnancy average	335	0.8	1.2	1.9	<LOD – 19	1.3 (2.0)	82%
<b>Neonatal TSH<sup>d</sup> analyses</b>							
Closest measurement	364	0.7	1.2	1.9	<LOD – 48	1.2 (2.3)	82%
Farthest measurement	364	0.7	1.1	1.8	<LOD – 27	1.1 (2.3)	82%
Pregnancy average	364	0.8	1.2	1.9	<LOD – 24	1.3 (2.0)	82%

<sup>a</sup> GSD = geometric standard deviation.

<sup>b</sup> Thyroid hormone.

<sup>c</sup> Limit of detection (LOD) = 0.4 $\mu\text{g/L}$ .

<sup>d</sup> Thyroid-stimulating hormone